



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2013

---

## Measuring threat status

Schmidt, B R

Abstract: What does it mean if a species is “endangered and red-listed”? Using the example of the 2005 Red List of Amphibians, this article aims to demonstrate how a species’ threat status is assessed.

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-85724>


Journal Article

Published Version

Originally published at:

Schmidt, B R (2013). Measuring threat status. Hotspot, 28:16.





# HOTSPOT



## Measuring biodiversity

Research and practice in dialogue  
Bulletin of the Swiss Biodiversity Forum

28 | 2013



# Authors

**Dr. Matthias Albrecht** is an ecologist and researcher with the Agricultural Landscapes and Biodiversity Group at the Agroscope Reckenholz-Tänikon Research Station. He participates in the QUESSA project under the 7th Framework Programme of the EU. He focuses on biodiversity and ecosystem services in agro-ecosystems and approaches to support these.

**Dr Ariel Bergamini** is a botanist and heads the Ecosystem Dynamics research group at the Swiss Federal Institute for Forest, Snow and Landscape Research WSL as well as the programme "Monitoring the Effectiveness of Habitat Conservation in Switzerland". Further interests include conservation biology with respect to flowering plants and mosses.

**Simon Birrer** heads the Förderung der Vogelwelt (Support for birdlife) department at the Swiss Ornithological Institute and focuses on applied projects in farming and forestry.

**Dr Stefan Eggenberg** studied plant taxonomy and vegetation ecology and trained as a scientific draftsman. Formerly co-owner of the Atelier für Naturschutz und Umweltfragen (UNA) in Bern, he is currently Director of Info Flora, the national documentation and information centre for Swiss flora.

**Dr Lisa Garnier** holds a PhD in ecology. As science journalist, writer and project coordinator, she specialises on conveying biodiversity issues to the wider public. She maintains the Vigie-Nature blog of the French National Natural History Museum in Paris and develops interactive scientific experiments.

**Christian Ginzler** is a biologist with the Swiss Federal Institute for Forest, Snow and Landscape Research WSL. He heads the Remote Sensing research group and primarily works on aerial photograph interpretation, photogrammetry and image analysis as tools to measure landscape change.

**Dr Yves Gonseth** heads the Swiss Biological Records Center (CSCF) where he is in charge of maintaining contacts with field researchers (primarily entomologists) and cantonal and federal authorities in the areas of species and habitat protection and with relevant international institutions.

**Anne-Laure Gourmand** works at the French National Natural History Museum in Paris, developing science programmes for Vigie-Nature and implementing them with local stakeholders. She coordinates STELI, a programme to monitor dragonfly populations in France.

**Dr Gabriela Hofer** is a biologist in the Agricultural Landscapes and Biodiversity Group at Agroscope Reckenholz-Tänikon Research Station. She develops approaches to illustrate species and habitat dynamics in open cultural landscapes and identify the contributions of ecological compensation areas to maintaining species diversity.

**Prof. Dr Rolf Holderegger** teaches at the Swiss Federal Institute of Technology and heads the Biodiversity and Conservation Biology research unit at the Swiss Federal Institute for Forest, Snow and Landscape Research WSL. He is responsible for overall administration of the programme "Monitoring the Effectiveness of Habitat Conservation in Switzerland".

**Dr Markus Jenny** is a biologist at the Swiss Ornithological Institute where he is in charge of agricultural projects at the interface of research, implementation, markets and policy. He presides over the Vision Landwirtschaft association, a think-tank of independent experts on farming matters.

**Dr Marc Kéry** is a population ecologist at the Swiss Ornithological Institute. His research interests include large-scale modelling of species distribution and abundance, population models, and modelling measurement error processes in ecological field studies.

**Dr Meinrad Küchler** works in the Ecosystem Dynamics research group at the Swiss Federal Institute for Forest, Snow and Landscape Research WSL. He focuses on statistical data analysis and modelling ecological change in different habitat types in Switzerland.

**Dr Enrique Lara** is a researcher at the University of Neuchâtel where he studies micro-eukaryotes (algae, fungi and a variety of unicellular organisms). He is particularly interested in their evolutionary history, ecology, geographic distribution and their enormous diversity.

**Dr Lukas Mathys** is a biologist acting as project manager at the Sigmaplan AG consultancy. In a variety of projects, he deals with both substantive and technical aspects of the collection, evaluation and communication of biodiversity data.

**Prof. Edward Mitchell** has been in charge of the Soil Biology Laboratory of the University of Neuchâtel since 2009 and has co-managed the Neuchâtel Botanic Garden since 2011. His interests include the ecology and biodiversity of soil organisms with a focus on protozoa.

**Dr Marco Moretti** is an ecologist and team leader at the Swiss Federal Institute for Forest, Snow and Landscape Research WSL in Bellinzona. For the past ten years he has studied aspects of biodiversity and biocoenoses and their relationships to ecosystem processes and services along a range of environmental gradients as well as under laboratory conditions.

**Prof. Jan Pawlowski** heads the Laboratory for Molecular Evolution and Ecology of Protists of the Department of Genetics and Evolution at Geneva University. His research explores the evolutionary history of eukaryotes. He also manages the Swiss Barcode of Life (SwissBOL) network.

**Dr Lukas Pfiffner** is an agro-ecologist managing biodiversity and conservation projects at the Research Institute of Organic Agriculture FiBL with a focus on ecological systems optimisation and trophic interactions of arthropods and soil organisms in a variety of cropping systems.

**Dr Benedikt Schmidt** works at karch – the coordination centre for amphibian and reptile conservation in Switzerland – and leads a research group at Zurich University. By combining research and practical work he contributes to evidence-based conservation.

**Dr Eva Spehn** is on the scientific staff of the Swiss Biodiversity Forum and manager of the international Global Mountain Biodiversity Assessment network which maintains an online portal for biodiversity data for the world's mountain regions ([www.mountainbiodiversity.ch](http://www.mountainbiodiversity.ch)). She is a member of the GBIF Swiss Commission and DIVERSITAS delegate to GBIF.

**Dr Sibylle Stöckli** is a project manager at the Research Institute of Organic Agriculture FiBL in the areas of biodiversity, climate change and ecosystem functions with a focus on entomology and crop protection.

**Silvia Stofer** heads the Biodiversity Assessment research group within the Biodiversity and Conservation Biology unit at the Swiss Federal Institute for Forest, Snow and Landscape Research WSL. Her responsibilities include maintaining the national data base of lichens in Switzerland (Swiss-Lichens).

**IMPRINT** The Swiss Biodiversity Forum promotes the exchange of knowledge between biodiversity research, administrations, practitioners, policy-makers and wider society. HOTSPOT is one of the tools facilitating this exchange. HOTSPOT is published twice a year in German, English and French. It is available for download at [www.biodiversity.ch](http://www.biodiversity.ch). HOTSPOT 29|2014 will be published in May 2014 and will address the topic of Energy and Biodiversity. **Publisher:** © Forum Biodiversität Schweiz, Bern, November 2013. **Editors:** Dr Gregor Klaus (gk), Dr Daniela Pauli (dp). **Translation into English:** Christopher Hay. **Graphic design:** Esther Schreier, Basel. **Photos:** Photo credits are listed with the captions. **Printing:** Print Media Works, Schopfheim im Wiesental. **Paper:** Circle matt 115 g/m<sup>2</sup>, 100% recycled. **Print edition:** 3,300 copies in

German, 1,100 copies in French, 1,000 copies in English. **Contact:** Forum Biodiversität Schweiz, Schwarztorstr. 9, CH-3007 Bern, Switzerland, Tel. +41 (0)31 312 02 75, [biodiversity@scnat.ch](mailto:biodiversity@scnat.ch), [www.-biodiversity.ch](http://www.-biodiversity.ch). **Managing Director:** Dr Daniela Pauli. **Cost of production:** 15 CHF/copy.

Sharing biodiversity knowledge with all those who are interested in these matters is important to us. We would therefore like to continue to offer HOTSPOT free of charge. Any financial support will be greatly appreciated. HOTSPOT donations account: PC 30-204040-6. Manuscripts are subject to editorial revision. Authors' contributions do not necessarily reflect the opinions of the Swiss Biodiversity Forum.

sc | nat 

Science and Policy  
Platform of the Swiss Academy of Sciences  
Swiss Biodiversity Forum

**Cover** (top to bottom):

1. Diverse micro-organisms (photo credit: Edward A. D. Mitchell); 2. Determining the diversity of fruit varieties (photo credit: ProSpecieRara Basel); 3. Archived butterfly diversity (photo credit: Beat Ernst Basel); 4. Biologists working in the field (photo credit: Edi Stöckli)

# Red lists

## Measuring threat status


Benedikt R. Schmidt, Koordinationsstelle für Amphibien- und Reptilienschutz in der Schweiz (karch), CH-2000 Neuchâtel, benedikt.schmidt@unine.ch

**What does it mean if a species is “endangered and red-listed”? Using the example of the 2005 Red List of Amphibians, this article aims to demonstrate how a species’ threat status is assessed.**

The contracting authority of the 2005 Red List of Amphibians, the Swiss Federal Office for the Environment FOEN, specified that the methodology used by the World Conservation Union (IUCN) should be applied. The IUCN has defined both red list categories as well as quantitative criteria which determine to which of the categories a species is assigned. An examination of the IUCN’s methodology showed that for amphibians native to Switzerland only the criteria “geographic range” and “declining population” can meaningfully be applied. As karch – Swiss Amphibian and Reptile Conservation Programme – maintains a distribution database which provides information on the number of populations of species of amphibians in Switzerland, and given that changes in the number of populations may be used to infer a population reduction, the “declining population” criterion appeared appropriate.

For all pond-dwelling species (except common species, namely the European common frog, European toad and Alpine newt), 25 breeding sites (i.e. populations) each were selected at random. As each of the breeding sites often hosts several species, the resulting sample sizes were in the order of 25 to 100 breeding sites per species. A total of 300 breeding sites of amphibians were sampled, each being visited four times by specialists. The aim was to record the amphibian species (still) present. Four visits were needed due to the species’ phenology; these multiple visits also allowed an estimate of species’ detection probability. The estimates of detection probability allowed calculation of the number of populations overlooked. It was shown that, thanks to the four visits to each site, practically no population remained undetected.

The calculations for the “declining population” criterion were based on the popu-



The map shows, for the Yellow-bellied toad (*Bombina variegata*), the results of field work undertaken to update the red list. Green dots denote past records which were confirmed, while gray dots denote past records which could not be confirmed. Red dots denote new populations. Copyright: karch and Swisstopo.

lation decline as determined from field survey data. For example, if a species previously had 100 populations in the overall sample and only 50 populations could be confirmed it was deemed to have suffered a 50% population decline.

It would also have been possible to record a population increase. In addition to breeding sites in which the species had been recorded in the past, sites were included in the sample at which the species had not been observed in the past, including some newly created wetlands. If a species was recorded at breeding sites where it had not previously been present, this would indicate a population increase. However, relatively few such populations were detected and mostly only concerned common species such as the European common frog or Alpine newt.

The “effective range” was determined by first calculating the number of current populations (number of populations in database multiplied by population decline)

and then determining the area inhabited by each of the populations. For frogs this area was taken to cover a circle with a radius of one kilometre while for newts an 0.5 kilometre radius was applied. The number of current populations multiplied by their individual range gives the species’ “effective range” in Switzerland.

For a reliable estimate of population decline and range size, and to assign the species to IUCN red list categories, it is crucial that the karch database held good records of species distribution in Switzerland, that random sampling of populations in the field was undertaken, and that the data were analysed using modern statistical methods (also see the article on p. 8).